

## Article

# Should all patients with generalised tonic-clonic seizures have their troponin levels measured?

Egiz, Abdullah Mohammed abousaleh ma, Nautiyal, Hritik, Al-Shaharli, Radhi Forat Radhi, Fazail, Abdurrahman Hazrin and Teo, Bervin

Available at <http://clock.uclan.ac.uk/37399/>

*Egiz, Abdullah Mohammed abousaleh ma ORCID: 0000-0003-0304-7982, Nautiyal, Hritik, Al-Shaharli, Radhi Forat Radhi, Fazail, Abdurrahman Hazrin and Teo, Bervin (2021) Should all patients with generalised tonic-clonic seizures have their troponin levels measured? British Journal of Hospital Medicine, 82 (4). ISSN 1750-8460*

It is advisable to refer to the publisher's version if you intend to cite from the work.  
10.12968/hmed.2020.0581

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

# British Journal of Hospital Medicine

## Should all patients with generalised tonic-clonic seizures have their troponin measured? A case report and review of literature.

--Manuscript Draft--

Manuscript Number:	hmed.2020.0581R1
Full Title:	Should all patients with generalised tonic-clonic seizures have their troponin measured? A case report and review of literature.
Short Title:	Cardiac Troponin T elevation in a patient following a generalised tonic-clonic seizure.
Article Type:	Case report
Keywords:	Seizures; Epilepsy; Tonic-Clonic; Troponin T; Myocardial Ischemia.
Corresponding Author:	Bervin Teo Chong Wei West Cumberland Hospital Whitehaven, Cumbria UNITED KINGDOM
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	West Cumberland Hospital
Corresponding Author's Secondary Institution:	
First Author:	Abdullah Egiz
First Author Secondary Information:	
Order of Authors:	Abdullah Egiz Hritik Nautiyal Radhi Al-Shaharli Abdurrahman Fazail Bervin Teo
Order of Authors Secondary Information:	
Abstract:	This report presents the case of a 76-year-old male patient who presented with two tonic-clonic seizures within two hours, each episode lasting approximately two minutes. The first seizure was not accompanied by tongue-biting or incontinence. However, upon admission to the emergency department, the second seizure episode was associated with tongue-biting, incontinence and upward-rolling of the eyes. The patient's national early warning score upon admission was 3. Physical examinations were unremarkable. Biochemistry results showed cardiac troponin T (cTnT) elevation at 52ng/L (<14 ng/L), with a further elevation to 57ng/L after 5 hours.
Suggested Reviewers:	
Response to Reviewers:	Dear Reviewer,  We hope that you are doing well during the current challenging times.  We are thankful for your constructive feedback and would like to inform you that we have made the recommended changes given by yourself. This includes reducing the word count of the main text below 800 words.  We look forward to hearing back from you soon.  Kind Regards, Bervin Teo
Additional Information:	

Question	Response
Please enter the word count of your manuscript, <b>excluding references and tables</b>	943

# **Should all patients with generalised tonic-clonic seizures have their troponin measured? A case report and review of literature.**

**Short title: Cardiac Troponin T elevation in a patient following a generalised tonic-clonic seizure.**

Abdullah Egiz, Medical student<sup>1</sup>  
Hritik Nautiyal, Medical student<sup>1</sup>  
Radhi Al-Shaharli, Medical student<sup>1</sup>  
Abdurrahman Fazail, Medical student<sup>1</sup>  
Bervin Teo, Locum Medical Consultant<sup>2</sup>

<sup>1</sup>School of Medicine, University of Central Lancashire

<sup>2</sup>West Cumberland Hospital, Whitehaven

Corresponding author: Bervin Teo, [bervin.teo2@nhs.net](mailto:bervin.teo2@nhs.net), +44 (0) 7753 458884

## **Abstract**

This report presents the case of a 76-year-old male patient who presented with two tonic-clonic seizures within two hours, each episode lasting approximately two minutes. The first seizure was not accompanied by tongue-biting or incontinence. However, upon admission to the emergency department, the second seizure episode was associated with tongue-biting, incontinence and upward-rolling of the eyes. The patient's national early warning score upon admission was 3. Physical examinations were unremarkable. Biochemistry results showed cardiac troponin T (cTnT) elevation at 52ng/L (<14 ng/L), with a further elevation to 57ng/L after 5 hours.

## **Keywords**

Seizures; Epilepsy; Tonic-Clonic; Troponin T; Myocardial Ischemia.

## **Acknowledgment**

We thank the patient for giving us the privilege and honour to share his case and learn from it.

## **Conflict of interest statement**

No, there are no competing interests for any author

## **Should all patients with generalised tonic-clonic seizures have their troponin measured? A case report and review of literature.**

### **Abstract**

This report presents the case of a 76-year-old male patient who presented with two tonic-clonic seizures within two hours, each episode lasting approximately two minutes. The first seizure was not accompanied by tongue-biting or incontinence. However, upon admission to the emergency department, the second seizure episode was associated with tongue-biting, incontinence and upward-rolling of the eyes. The patient's national early warning score upon admission was 3. Physical examinations were unremarkable. Biochemistry results showed cardiac troponin T (cTnT) elevation at 52ng/L (<14 ng/L), with a further elevation to 57ng/L after 5 hours.

### **Case report**

A 76-year-old male brought in by ambulance to the emergency department due to an episode of nocturnal generalised tonic-clonic seizure (GTCS). Collateral history from the patient's wife revealed the seizure lasted for two minutes and it was associated with generalised stiffness of head, body and limbs with arched back followed by clonic jerks of head, body, limbs and upward eye-rolling. However, within two hours post-admission the patient experienced a second GTCS associated with tongue-biting, backward eye-rolling and incontinence, lasting for two minutes. Another history was then taken from the patient once he recovered. The patient denied antecedent symptoms such as chest pain, shortness of breath, fever, neck stiffness or visual changes. Moreover, there was no history nor symptoms of focal neurological deficit or cranial nerve involvement. The patient has a past medical history of unprovoked first-onset nocturnal seizure 4 months before this episode and has no known history of cardiac or renal diseases. Before admission, he was taking Fostair (200/6) and salbutamol (PRN).

On examination, the patient was alert and tachycardic with heart rate of 128 bpm. He had an oxygen saturation level of 94% on room air. Other vital signs and physical examinations were otherwise unremarkable. National early warning score (NEWS) was 3. His postictal Glasgow Coma Score (GCS) was 15/15.

An electrocardiogram (ECG) revealed voltage criteria for mild left ventricular hypertrophy (LVH). Baseline blood samples were obtained and cTnT levels were also requested to rule out myocardial infarction as the potential source of the syncope. Biochemistry results showed an elevation of lactate at 13mmol/L (0.1-1mmol/L) and a low bicarbonate at 16mmol/L (22-26 mmol/L) along with a cTnT elevation at 52 ng/L (<14 ng/L), with a further elevation to 57ng/L after 5 hours. White cell count (WCC) was slightly elevated at  $16.1 \times 10^9/L$  ( $4-11 \times 10^9/L$ ) with neutrophils elevated at  $10.8 \times 10^9/L$  ( $2.0-7.5 \times 10^9/L$ ). Blood sugar levels were normal. Computed tomography of the brain did not reveal any cerebrovascular disease or space-occupying lesion.

The patient received Intravenous fluid therapy of 500 mL, 0.9% saline for fluid resuscitation. Subsequently, intravenous diazepam was administered at a rate of 5 mg per minute. 250 mg of Levetiracetam was prescribed to be taken orally twice daily. 625 mg of co-amoxiclav was administered as a precautionary empirical broad-spectrum antibiotic considering the raised neutrophils, WCC and lactate. The patient

responded well to the treatment and remained stable. He was discharged in two days and booked in for a neurological assessment in 2 weeks.

## **Discussion**

Patients with GTC seizures who have elevated postictal cardiac troponin levels upon biochemical investigation are either elderly, have cardiac risk factors or both (Chatzikonstantinou et al., 2015). This can be highlighted by the fact that epilepsy studies involving young patients and with mean cohort ages of 21 and 44 consistently report no troponin elevation following seizures (Eskandarian et al., 2011; Hajsadeghi et al., 2009). In contrast, the study by Chatzikonstantinou et al (2015) which included a higher mean cohort age of 59 years, reported elevations in troponin levels following focal and generalized seizures. In addition, a systematic review by Sieweke et al. (2012), links cardiovascular ischemia as a contributing factor to seizure-related elevations in cTnT. Hence, our case is consistent with the findings of these studies as it reports troponin elevation postictally in a 76-year-old male patient.

There is limited research that investigates the link between seizures and troponin rise and therefore the exact pathophysiology is controversial (Fawaz et al., 2014). However, considering the autonomic innervation of the heart, periseizure cardiac changes are not surprising according to Jansen and Lagae (2010). It is postulated that the hyperadrenergic state during convulsions leads to a perfusion/demand mismatch resulting in myocardial injury and rise in troponins (Gaspard, 2019). Although troponins are considered to be specific to cardiac injury, troponin elevation has been reported in non-acute coronary syndrome events such as myocarditis, pericarditis, sepsis and rhabdomyolysis (Fawaz et al., 2014). As our patient did not have any of the above conditions, the possibility of a minor ischaemic myocardial injury due to the perfusion/demand mismatch seems more likely. It is valid to assume that the elderly are more prone to perfusion/demand mismatch injuries considering the age-related vascular changes that lead to greater myocardial demand (Fawaz et al., 2014).

We conclude through this case report that a GTCS cannot be assumed to be an innocent event for the patient's heart, especially in older age groups. A comprehensive postictal cardiovascular assessment is therefore warranted. More accurate imaging modalities such as contrast-enhanced cardiovascular MRI for assessing subendocardial damage can be employed post-GTCS in patients with epilepsy to further investigate the link between cardiac injury and seizures.

## **Learning points**

- GTC seizures seem to prompt the release of cTnT with predominance in patients with cardiovascular risk factors, especially in advanced age. This points towards the hypothesis that myocardial injury during the peri-seizure period causes the rise in cardiac troponins.
- A comprehensive postictal cardiovascular assessment is warranted including, but not limited to, an ECG, echocardiogram and troponins assay.
- Low clinical suspicion and lack of effort to explore for myocardial injury and other medical comorbidities post-GTC seizure could miss the diagnosis of potential co-existing conditions.

## **Key words**

Seizures; Epilepsy; Tonic-Clonic; Troponin T; Myocardial Ischemia.

## References

- Chatzikonstantinou, A., Ebert, A.D., Hennerici, M.G., 2015. Temporal seizure focus and status epilepticus are associated with high-sensitive troponin I elevation after epileptic seizures. *Epilepsy Research* 115, 77–80.  
<https://doi.org/https://doi.org/10.1016/j.eplepsyres.2015.05.013>
- Eskandarian, R., Asghari, N., Darban, M., Ghorbani, R., 2011. Cardiac Troponin Levels Following Complicated and Uncomplicated Epileptic Seizures. *Archives of Medical Research* 42, 439–442. <https://doi.org/https://doi.org/10.1016/j.arcmed.2011.09.002>
- Fawaz, A., Nasreddine, W., Makke, Y., Atweh, S., Wazne, J., Arabi, M., Beydoun, A., 2014. Association of cardiovascular risk factors and troponin elevation after generalized tonic-clonic seizures. *Seizure - European Journal of Epilepsy* 23, 146–150.  
<https://doi.org/10.1016/j.seizure.2013.11.006>
- Gaspard, N., 2019. Heartbreakers-Cardiac Stress After Uncomplicated Generalized Convulsive Seizures. *Epilepsy currents* 19, 246–248.  
<https://doi.org/10.1177/1535759719856592>
- Hajsadeghi, S., Afsharian, S., Fereshtehnejad, S.-M., Keramati, M.R., Mollahoseini, R., 2009. Serum levels of cardiac troponin I in patients with uncomplicated epileptic seizure. *Archives of Medical Research* 40, 24–28.  
<https://doi.org/10.1016/j.arcmed.2008.10.002>
- Jansen, K. & Lagae, L. 2010, Cardiac changes in epilepsy. *Seizure - European Journal of Epilepsy*, vol. 19, no. 8, pp. 455-460. <https://doi.org/10.1016/j.seizure.2010.07.008>
- Sieweke, N., Allendörfer, J., Franzen, W., Feustel, A., Reichenberger, F., Pabst, W., Krämer, H.H., Kaps, M. & Tanislav, C. 2012. Cardiac Troponin I elevation after epileptic seizure. *BMC Neurology*, vol. 12, no. 1, pp. 58. <https://doi.org/10.1186/1471-2377-12-58>